

FINAL REPORT
NAGW-1081
SPECTRAL REFLECTANCE OF PLANETARY SURFACES

Principal Investigator: John B. Adams

University of Washington

30 May 1991

This report emphasizes the final year of the grant period. Earlier work, dating back to 1981 is summarized in the publications that are cited below.

Detectability Analysis. An overall theme of our research has been to evaluate and interpret spectroscopic evidence for compositional variability on Mars. We have assessed the compositional variability that could be detected spectroscopically under computer-simulated "best case" conditions using all available wavelength measurements. We used laboratory spectra of Martian analog materials including basalt, weathered basaltic palagonite and other mineral and rock types. Laboratory spectra were convolved with Viking Lander and Viking Orbiter spectral channels and the resulting pseudospectra were analyzed for detectability. In the detectability analysis the spectrum of one material is evaluated relative to a specified background which may consist of one or more spectrally distinct materials. The detectability of the "target" material depends on the spectral contrast with the background and on system parameters, including the number and wavelength of the camera channels and the signal-to-noise ratio of the measurements.

Simulations also were made of telescopic spectra by normalizing laboratory spectra at 0.56um. Detectability thresholds were then determined for various target materials and backgrounds. Thresholds were measured at different signal-to-noise levels for laboratory measurements and for the simulated Viking Lander, Viking Orbiter, and telescopic data. Our results indicate that there is a wide range of detectability thresholds for Martian analog materials and that, in general, detectability of most materials is poor for systems with few channels such as Viking Orbiter, and that it also is poor for normalized data such as the telescopic measurements.

An example is the detectability threshold of andesitic ash (Mt. St. Helens) in a background of basalt and weathered basaltic palagonite. We posed the question: how much andesitic ash would have to be present to be spectrally detectable in the presence of basalt and palagonite? Alternatively, if the present spectroscopic and other evidence is correct that the surface of Mars is dominated by basalts and weathered palagonitic dust, what other materials (such as andesite) might be present that might be spectrally indistinguishable from the two main materials and their mixtures? We showed that andesitic ash is detectable using laboratory spectra at amounts of a few percent. However, when measured by Viking Orbiter the ash must be a major component of the

(NASA-CR-194310) SPECTRAL
REFLECTANCE OF PLANETARY SURFACES
Final Report (Washington Univ.)
7 p

N94-70603

Unclass

surface to be detected. Similarly, to be detected in the telescopic spectra with a signal-to-noise ratio of 50:1 the andesitic ash would have to comprise 15 to 20% of the surface.

The poor detectability of andesitic ash in the telescopic simulation occurs in spite of high spectral resolution. The problem lies in the normalization of the data which eliminates the "albedo" information. Most researchers analyzing telescopic spectra today work with normalized data because they are interested primarily in identifying specific absorption bands that can be assigned to minerals. Unfortunately, most absorption bands in the Martian spectra are weak (excepting the Fe³⁺ ones) and it is difficult to isolate weak mineral bands from atmospheric ones and from the instrumental noise. Current debates about scapolite illustrate the point.

Telescopic measurements are not being used to full advantage. Our modeling results indicate that if the albedo information could be recovered from the telescopic data the detectability of all materials would be increased substantially. For example, one then could examine whether a large fraction of scapolite would be detectable using the measurements over the rest of the spectrum, outside of the absorption bands. We are aware of the reasons for making normalized telescopic measurements, and of the difficulties in acquiring absolute data. However, the effort may be worthwhile.

Telescopic CCD Images. Beginning in 1988 and continuing through 1990 a large number of high spectral resolution CCD images were acquired at Mauna Kea and Pic du Midi. Some of these data have been published and some are being reduced. Although of very high spectral resolution, these measurements have been difficult to calibrate and, therefore, have been difficult to compare and to interpret in terms of laboratory reference spectra. This fundamental problem of calibrating spectral measurements that are obtained by various instruments under different illumination and atmospheric conditions has been one focus of our research for several years.

Part of our research effort in the past year has been to calibrate and interpret high spectral resolution telescopic spectra and CCD images in collaboration with T. McCord, J. Bell, and P. Pinet. In working with the telescopic CCD data we have addressed one particularly difficult problem, namely, that spectra are affected by the rotation of Mars during and between measurements. When spectral measurements are sequenced in time the field of view shifts as a function of wavelength. Smearing during measurement at a single wavelength increases the instantaneous field of view; however, movement during multiple (wavelength) measurements poses a difficulty in registering the multispectral data into an image cube. Registration can be approximated by shifting pixel arrays by a time-distance function, but accurate registration requires tie- points on the surface. Unfortunately, surface features vary in contrast with wavelength, and cannot be counted on for registration.

Terrestrial analogs. We have been working with J. Bell (U. of Hawaii and our laboratory) and R. Morris (JSC) to further evaluate the spectra of the bright areas of Mars in the visible and near-infrared. We have found a new sample of Hawaiian palagonitic tephra that has the best fit that we have seen so far to the telescopic spectra of martian dust. The sample occurs at the contact between a basalt dike and a basalt tephra cone. We have proposed that the palagonitic sample formed under conditions that also could have occurred on the surface of Mars as a result of volcanic or impact thermal pulses applied to basaltic tephra containing adsorbed water or ice. We have examined the magnetic and Mossbauer properties of the iron phases in the sample, and have evaluated the reflectance spectra in terms of the iron oxide / oxyhydroxide mineralogy.

PUBLICATIONS 1981 - 1990

- McCord T.B., Clark R.N., McFadden L.A., Pieters C.M., and Adams J.B., 1981 Moon: Near-infrared spectral reflectance, a first good look, J. Geophys. Res. **86**, 10883-10892.
- Adams J.B., Pieters C.M., Metzger A.E., Adler I., McCord T.B., Chapman C.R., Johnson T.V., and Bielefeld M.J., Remote sensing of basalts in the solar system in: Basaltic Volcanism on the Terrestrial Planets, chapter 2, 440-485. Pergamon 1981.
- Evans D.L. and Adams J.B., 1981 Comparison of spectral reflectance properties of terrestrial and martian surfaces determined from LANDSAT and Viking Orbiter multispectral images (abs.), Lunar and Planetary Sci. XI, Part 1, 271-273.
- Evans D.L., Farr T.G., and Adams J.B., 1981 Spectral reflectance of weathered terrestrial and martian surfaces, Proc. Twelfth Lunar and Planet. Sci. Conf., 1473-1479.
- Gaddis L.R., Adams J.B., Hawke B.R., Head J.W., McCord T.B., Pieters C.M., and Zisk S., 1981 Characterization and distribution of pyroclastic units in the Rima Bode region of the moon (abs.), Lunar and Planetary Sci. XII, Part 1, 318-320.
- McCord T.B., Clark R.N., Hawke B.R., McFadden L.A., Owensby P.D., Pieters C.M., and Adams J.B., 1981 Remote detection of olivine, pyroxene, and plagioclase: Analysis of three lunar sites (abs.), Lunar and Planetary Sci. XII, Part 2, 679-681.
- Pieters C.M., Adams J.B., and McCord T.B., 1981 Copernicus' peak: Possible cumulate olivine (abs.), Lunar and Planetary Sci. XII, Part 2, 833-835.
- Evans D.L., and Adams J.B., 1981 Derivation of compositional information from multispectral images (abs.), Int. Geologic. and Remote Sensing Symp., Wash D.C.
- Evans, D.L., Adams, J.B., Farr, T.G., and Roush, T.L., 1981 Terrestrial and synthetic analogs to Martian weathering products (abs.), Third World Mars Conf., Pasadena.
- Adams J.B., Evans D.L., Farr T.G., Smith M.O., Staley J.T., Palmer F., Borns D.J., Curtiss B., Taylor-George S., Roush T.L., and Bjorkman E., 1982 Interpretation of weathered surfaces in arid regions using LANDSAT multispectral images, Proceedings of conference: Remote Sensing of Arid and Semi-arid Lands, Vol. II, Cairo, Egypt.

- McCord T., El Baz F., and Adams J.B., 1982 The nature and extent of erosional and depositional features and rock and soil units in the Kharga Oasis region, Egypt, as determined from remote sensing (abs.), Conf. Remote Sensing of Arid and Semi-arid Lands, Cairo, Egypt.
- Pieters C., Adams J.B., Head J., McCord T., and Zisk S., 1982 Primary ejecta in crater rays: The Copernicus example (abs.), Lunar and Planetary Sci. XIII Part 2, 623-624.
- Evans D.L., Farr T.G., Williams S., Smith M.O., and Adams J.B., 1982 Analysis of multisensor image data set of Death Valley, California (abs.), Geol. Soc. Amer. **14**, 4, 162-163.
- McCord T.B., Singer R.B., Hawke B.R., Adams J.B., Evans D.L., Head J.W., Mouginis-Mark P.J., Pieters C.M., Huguenin R.L., and Zisk S.H., 1982 Mars: Definition and characterization of global surface units with emphasis on composition, J. Geophys. Res. **87**, B12, 10129-10148.
- Johnson P.E., Smith M.O., Taylor-George S., and Adams J.B., 1982 A semi-empirical method for analysis of the reflectance spectra of binary mineral mixtures, J. Geophys. Res. **88**, B4, 3557-3561.
- Farr T. and Adams J.B., 1984 Rock coatings in Hawaii, Geol. Soc. Amer. Bull. **95**, 1077-1083.
- Smith M.O., Johnson P.E., and Adams J.B., 1984 A strategy for interpretation of multispectral reflectance data from planetary surfaces: Methodolgy (abs.), Lunar and Planetary Science XV, 798-799.
- Johnson P.E., Smith M.O., and Adams J.B., 1984 Application of a strategy for interpretation of multispectral reflectance data from planetary surfaces (abs.), Lunar and Planetary Science XV, 407-408.
- Curtiss B., Adams J.B., and Ghiorso M., 1985 Origin, development and chemistry of silica-alumina rock coatings from the semi-arid regions of the island of Hawaii, Geochim. et Cosmochim. Acta **40**, 49-56.
- Smith M.O., Johnson P.E., and Adams J.B., 1985 Quantitative determination of mineral types and abundances from reflectance spectra using principal components analysis, Proc. Lunar and Planetary Sci. Conf. XV, J. Geophys. Res. **90**, C797-C804.
- Johnson P.E., Smith M.O., and Adams J.B., 1985 Quantitative analysis of planetary reflectance spectra with principal components analysis, Proc. Lunar and Planetary Sci. Conf. XV, J. Geophys. Res. **90**, C805-C810.
- Curtiss B.E., Smith M.O., and Adams J.B., 1985 Separation of hydrothermal Fe (III) oxyhydroxides from weathering produced Fe (III) oxyhydroxides in multispectral images (abs.), Remote Sensing for Exploration Geology, Fourth Thematic Conf. ERIM, 89.
- Adams J.B., Smith M.O., and Johnson P.E., 1985 Viking Lander 1: A new map of rock and soil types (abs.), Lunar and Planetary Science XVI, part 1, 3-4.
- Smith M.O. and Adams J.B. Determining mineral types and abundances from reflectance measurements, Reports of Planetary Geology Program 1984, NASA Technical Memorandum TM 87563, April 1985.
- Smith M.O. and Adams J.B., 1985 Interpretation of AIS images of Cuprite, Nevada using constraints of spectral mixtures, Proceedings of the Airborne Imaging Spectrometer Data Analysis Workshop, JPL Pub. **85-41**, 62-68.

- Smith M.O. and Adams J.B., 1985 Strategies for analysing mixed pixels in remotely sensed imagery, NASA/JPL Aircraft SAR Workshop Proceedings, JPL Pub. 85-39, 47-48.
- Pieters C.M., Adams J.B., Mouginis-Mark P.J., Zisk S.H., Smith M.O., Head J.W., and McCord T.B., 1985 The nature of crater rays: The Copernicus example, J. Geophys. Res. **90**, 12393-12413.
- Adams J.B., Smith M.O., and Johnson P.E., 1986 Spectral mixture modeling: A new analysis of rock and soil types at the Viking Lander I Site, J. Geophys. Res. **91**, 8098-8112.
- Smith M.O., Adams J.B., and Johnson P.E., 1986 Separation of effects of composition, illumination geometry, and calibration in Viking 1 and 2 Multispectral Images (abs.), Lunar and Planetary Sci. XVII, 807-808.
- Johnson P.E., Smith M.O., and Adams J.B., 1986 Quantitative spectral mixture modeling applied to Viking Lander images (abs.), Lunar and Planetary Sci. XVII, 398-399.
- Cooke S.S., Adams J.B., and Smith M.O., 1986 The use of spectral endmembers in geobotanical image interpretation (abs.), Fifth Thematic Conf., Remote Sensing for Exploration Geology, ERIM.
- Smith M.O., Adams J.B., Guinness E.A., and Arvidson R.E., 1987 Viking Orbiter multispectral images linked to lander images and laboratory analogs (abs.), Lunar and Planetary Science XVIII, 934-935.
- Shipman H. and Adams J.B., 1987 Detectability of minerals on desert alluvial fans using reflectance spectra, J. Geophys. Res. **92**, 10391-10402.
- Adams J.B., Smith M.O., Arvidson R.E., Dale-Bannister M., Guinness E.A., and Singer R., 1987 Surface composition of Mars: A Viking multispectral view, Mars: Evolution of Volcanism, Tectonism, and Volatiles, LPI Publ., Dec. 1987.
- Adams J.B. and Smith M.O., 1987 Spectral mixture modeling: Further analysis of rock and soil types at the Viking Lander sites, NASA TM.
- Smith M.O. and Adams J.B., 1988 Telescopic spectra of Mars linked to Viking Lander multispectral images and laboratory analogs (abs.), Lunar and Planetary Science XIX.
- Ming D.W., Morris R.V., Bernhard R.P., and Adams J.B., 1988 Mineralogy of a suite of Martian analog soils from Hawaii (abs.), Lunar and Planetary Science XIX.
- Smith M.O., Adams J.B., and Gillespie A.R., 1988 Evaluation and calibration of AVIRIS test-flight data: Owens Valley, CA, Final Report NASA Contract No. NAGW 1135.
- Arvidson R.E., Guinness E.A., Dale-Bannister M.A., Adams J.B., Smith M.O., Christensen P.R., and Singer R.B., 1989 Nature and distribution of surficial deposits in Chryse Planitia and vicinity, Mars, J. Geophys. Res. **94**, 1573-1587.
- Adams J.B., Smith M.O., and Gillespie A.R., 1989 Simple models for complex natural surfaces: A strategy for the hyperspectral era of remote sensing, Proc. IEEE Int. Geosci. and Remote Sensing Symp'89 I, 16-21.
- Smith M.O. and Adams J.B., 1989 Isolation of compositional variance from Viking Lander multispectral images (abs.), Lunar and Planetary Science Conference XX 3, 1024-1025.

Blount G., Greeley P.R., Christensen N., Lancaster N., Adams J.B., and Smith M.O., 1989 Determination of regional surficial geology for aeolian terrains utilizing a linear mixing algorithm for remote sensing data (abs.), NASA TM 4130, 255-256.

Adams J.B. and Smith M.O., 1990 Limits on the compositional variability of the Martian surface (abs.), Lunar and Planetary Science Conference XXI, 1-2.

Sabol D.E., Adams J.B., and Smith M.O., 1990 Predicting the spectral detectability of surface materials using spectral mixture analysis, Int. Geosci. and Remote Sensing Symp '90, (IGARSS'90), 2, 967-970.

Blount G., Smith M.O., Adams J.B., and Greeley R., 1990 Regional aeolian dynamics and sand mixing in the Gran Desierto: evidence from Landsat Thematic Mapper images, J. Geophys. Res., 95, 15463-15482.

Gillespie A.R., Smith M.O., Adams J.B., and Willis S.C., 1990 Spectral mixture analysis of multispectral thermal infrared images; Proc. Second Thermal Infrared Multispectral Scanner (TIMS) Workshop, JPL Pub. 90-95, 57-74.